

# Association of Prenatal Vascular Disruptions With Decreased Maternal Age

Mark S. Lubinsky\*

<sup>1</sup>Division of Genetics, Department of Pediatrics, Medical College of Wisconsin, Milwaukee

<sup>2</sup>Genetics Center, Children's Hospital of Wisconsin, Milwaukee

**Disruptions of fetal structures can create a variety of congenital anomalies. Some apparent prenatal vascular disruptions associate strongly with decreased maternal age, and are rare with older mothers. This is well-documented for gastroschisis, but similar findings with hydranencephaly suggest a general phenomenon that may also involve porencephaly, septo-optic dysplasia, early body stalk disruptions, certain hemifacial anomalies, and other findings. Prenatal vascular disruption may be a common cause of congenital anomalies, but its nature is unknown, and obvious environmental confounders associated with decreased maternal age may have only relatively small contributions. A protective effect for pregnancies of older mothers also remains a possibility. Am. J. Med. Genet. 69:237–239, 1997.**

© 1997 Wiley-Liss, Inc.

**KEY WORDS:** gastroschisis; hydranencephaly; maternal age; septo-optic dysplasia; vascular disruption

## INTRODUCTION

Prenatal vascular events can disrupt fetal structures, creating a variety of congenital anomalies [Van Allen, 1981]. Gastroschisis, one such apparent finding, shows a distinct epidemiology with a strong association with decreased maternal age (DMA) and a paucity of older mothers. Hydranencephaly, another such finding, shows a similar effect, and analysis of septo-optic dysplasia suggests a general phenomenon of unknown cause. Other anomalies of this sort may also be involved, although exceptions do exist. This may be a relatively common cause of a variety of congenital anomalies.

\*Correspondence to: Mark Lubinsky, M.D., Genetics Center, Children's Hospital of Wisconsin, P.O. Box 1997, Mail Station 716, Milwaukee, WI 53201.

Received 14 February 1995; Accepted 31 October 1995

## EVIDENCE FOR AN ASSOCIATION OF DMA WITH PRENATAL VASCULAR DISRUPTIONS

Gastroschisis, a ventral body wall defect, is a probable vascular disruption. It is the easiest of such anomalies to study in terms of frequency, ease of diagnosis, and clinical homogeneity. Multiple studies have shown a distinct epidemiology with a clear DMA effect and a paucity of older mothers [Curry et al., 1993].

For institutionalized patients with hydranencephaly, a vascular disruption of the brain [Friede, 1989; Hunter, 1993], 10/16 mothers were teenaged, significantly more than population and institutional controls. Only 1 mother was over 29 years old [Lubinsky et al., 1997]. In a population study, teenage mothers had an almost 10-fold greater risk than mothers over age 30 years [Lubinsky and Torfs, 1996].

Increased illegitimacy with hydranencephaly [Opitz et al., 1982; Friede, 1989; Lubinsky et al., 1997] may be secondary to a high rate of out-of-wedlock births to younger mothers.

Increased illegitimacy was also seen with porencephaly [Dekaban, 1965], which can be part of the same spectrum of brain vascular disruptions as hydranencephaly [Friede, 1989].

Younger mothers predominate with septo-optic dysplasia [Lippe et al., 1979]. This variable combination of septum pellucidum, optic nerve, and pituitary anomalies is almost always sporadic. It associates with porencephaly, a vascular anomaly, and makes little embryologic sense. Components develop at different times and arise from different tissues and processes. Vascular disruptions, perhaps involving the proximal trunk of the anterior cerebral artery, are more likely than a dysplasia or some other classical developmental process [Lubinsky, 1997].

## DISCUSSION

Gastroschisis, hydranencephaly, and septo-optic dysplasia all have significant excesses of younger mothers. The magnitude of this effect, plus a relative rarity of older mothers for at least the first two conditions [Curry et al., 1993; Lubinsky et al., 1997; Lubinsky and Torfs, 1996], differs greatly from patterns seen with most other congenital anomalies [Croen and Shaw, 1995], and suggests a common pathogenesis.

While alternatives have been proposed for vascular disruptive involvement with gastroschisis and septo-optic dysplasia, there are strong arguments for this process [Curry et al., 1993; Lubinsky, 1997]. However, there is little question for hydranencephaly [Hunter, 1993], and the distinct DMA epidemiology suggests a common link that is obscure with other proposed origins.

DMA effects can be complex. Even genetic traits can be affected (e.g., polydactyly and piebaldism in guinea pigs [Wright, 1926]), and the nature of the suggested phenomenon is unclear. It also should be noted that, despite a focus on risk factors at lower ages, a protective effect for older mothers could give a similar epidemiology even with a diverse set of causes.

Vascular disruptions are, indeed, diverse: diabetes, prenatal infections, embolization or coagulopathy from a deceased twin, including fetal papyraceus, twin-twin transfusions, and anatomic variants of the umbilical cord have all been implicated [Benirschke and Kaufmann, 1990; Hoyme et al., 1982], and clearly vascular findings can also be part of chromosomal or genetic syndromes at times (e.g., cutis aplasia [Manino et al., 1977; Der Kaloustian, 1993]).

However, most such etiologies should not show a DMA effect, and a paucity of older mothers suggests a single age-sensitive effect for most cases of gastroschisis and hydranencephaly. This may reflect at least relative causal homogeneity for these conditions, which are mostly sporadic, and rare with aneuploidy, Mendelian disorders, and nondisruption anomalies. Still, this is probably not absolute, and gastroschisis with multiple vascular anomalies does not appear to be age-related [Calzolari et al., 1995; Yang et al., 1992].

Greater heterogeneity of other disruption anomalies may obscure DMA effects, but possibilities exist with other cases. For body stalk agenesis and amelia with other gross body wall defects, all possible vascular disruptions [Luebke et al., 1990; Mastroiacovo et al., 1992] found a high rate of young mothers. Also, for cutis aplasia, Cockayne [1933] noted a tendency for affected children to be firstborn.

Hemifacial findings may also have vascular origins [Poswillo, 1973; Robinson et al., 1990]. In a California clinic, I found an average maternal age 3 years lower for isolated cases than cleft lip  $\pm$  palate controls, and no mothers over age 29 years. Numbers were small (12) but results were suggestive ( $P = 0.065$ ). Such analyses may help distinguish cases with a "standard" vascular disruptive etiology from those with other origins, such as hemifacial anomalies with epibulbar dermoids, which are difficult to see as vascular findings.

However, a DMA effect was absent with limb defects [Mastroiacovo and Botto, 1994] that are probably of vascular origin [Bavinck and Weaver, 1986]. Interestingly, this is the only vascular finding noted with chorionic villus sampling (cvs), which is often done for advanced maternal age [Mastroiacovo and Botto, 1994]. If this is a unique exception, then other disruptions might be potentiated with cvs at younger ages, giving anomalies not seen in current studies.

Confounding factors are possible, but unclear [Drongowski et al., 1991; Goldbaum et al., 1990; Haddow et al., 1993; Torfs et al., 1994; Werler et al., 1992a,b]. Low parity is not the answer alone [Hemminki et al., 1982]. High illegitimacy with por- and hydranencephaly [Opitz et al., 1982; Dekaban, 1965; Friede, 1989] may only be secondary [Lubinsky et al., 1997], but events such as attempted terminations are possible [Warkany et al., 1981]. Influences may also have changed: all cited illegitimate births were before 1985 and, despite some increases in gastroschisis rates [Curry et al., 1993], no epidemic of disruptions has been noted since [Martin et al., 1992], despite rising teenage illegitimacy [Guhleman, 1993].

Younger mothers also tend to have a lower socioeconomic status, but a dearth of older mothers with gastroschisis and hydranencephaly is hard to explain on this basis. Also, despite a higher rate of younger mothers and a generally lower socioeconomic status, gastroschisis is relatively rare, and a DMA effect smaller, in blacks than in whites [Center for Disease Control, 1993; Yang et al., 1992; Lubinsky et al., 1995].

Cocaine, with its vascular effects, might have a role, but pregnant users tend to be slightly older than other childbearing women, and greater use by blacks [Robins and Mills, 1993] is not reflected in racial incidences of gastroschisis [Center for Disease Control, 1993; Lubinsky et al., 1995; Yang et al., 1992]. Also, DMA observations in gastroschisis predate current drug use patterns, and no epidemic of disruptions has been noted recently [Martin et al., 1992], making a primary role unlikely. The DMA effect also suggests caution in studying cocaine teratogenesis: if vascular pathogenesises are being considered, then incidence rates should be from children of age-controlled mothers, and not from general population figures.

## REFERENCES

- Bavinck JNB, Weaver D (1986): Subclavian artery supply disruption sequence: Hypothesis of a vascular etiology for Poland, Klippel-Feil, and Mobius anomalies. *Am J Med Genet* 23:903-918.
- Benirschke K, Kaufmann P (1990): "Pathology of the Human Placenta," 2nd ed. New York: Springer-Verlag.
- Calzolari E, Bianchi F, Dolk H, Milan M, EUROCAT Working Group (1995): Omphalocele and gastroschisis in Europe: A survey of 3 million births, 1989-1990. *Am J Med Genet* 58:187-194.
- Centers for Disease Control (1993): Congenital malformations surveillance. Reprint, U.S. Department of Health and Human Services. *Teratology* 48:545-709.
- Cockayne EA (1933): "Inherited Abnormalities of the Skin." London: Oxford University Press, p 316.
- Croen LA, Shaw GM (1995): Young maternal age and congenital malformations: A population-based study. *Am J Public Health* 85:710-713.
- Curry CJR, Honore L, Boyd E (1993): The ventral wall of the trunk. In Stevenson RE, Hall JG, Goodman RM (eds): "Human Malformations and Related Anomalies," Vol II. New York: Oxford University Press, pp 869-905.
- Dekaban A (1965): Large defects in the cerebral hemispheres associated with cortical dysgenesis. *J Neuropathol Exp Neurol* 24:512-530.
- Der Kaloustian VM (1993): Skin. In Stevenson RE, Hall JG, Goodman RM (eds): "Human Malformations and Related Anomalies," Vol II. New York: Oxford University Press, pp 931-997.
- Drongowski RA, Smith RK Jr, Coran AG, Klein MD (1991): Contribution of demographic and environmental factors to the etiology of gastroschisis: A hypothesis. *Fetal Diagn Ther* 6:14-27.

- Friede RL (1989): "Developmental Neuropathology," 2nd ed. New York: Springer-Verlag, p. 322.
- Goldbaum G, Daling J, Milham S (1990): Risk factors for gastroschisis. *Teratology* 42:397-403.
- Guhleman P (1993): "Births to Wisconsin Teens: Trends 1960-1991," Vol 7. Health Data Review. Wisconsin DHSS, Center for Health Statistics, p 5.
- Haddow JE, Palomaki GE, Holman MS (1993): Young maternal age and smoking during pregnancy as risk factors for gastroschisis. *Teratology* 47:225-228.
- Hemminki K, Saloniemi I, Kyyronen P, Kekomaki M (1982): Gastroschisis and omphalocele in Finland in the 1970s: Prevalence at birth and its correlates. *J Epidemiol Community Health* 36:287-293.
- Hoyme HE, Jones KL, Van Allen MI, Saunders BS, Bic mechanisms in a population referred to a community-based stillbirth assessment program. *Am J Med Genet* 36:56-72.
- Hunter AGW (1993): Brain. In Stevenson RE, Hall JG, Goodman RM (eds): "Human Malformations and Related Anomalies," Vol II. New York: Oxford University Press, pp 1-107.
- Lippe B, Kaplan SA, La Franche S (1979): Septo-optic dysplasia and maternal age. *Lancet* 8133:92-93.
- Lubinsky MS (1997): Hypothesis: Septo-optic dysplasia is a vascular disruption sequence. *Am J Med Genet* 69:235-236.
- Lubinsky MS, Torfs C (1996): Decreased maternal age effect for isolated hydranencephaly in a population based study. Abstract: XVI David W. Smith Workshop on Malformations and Morphogenesis. Big Sky, Montana. In: Proceedings of the Greenwood Genetic Center, Vol 15, p 148.
- Lubinsky MS, Torfs C, Bateson T (1995): Lower age specific rates for gastroschisis in California blacks than in whites: An unexpected relationship of race and decreased maternal age effects in a population based study. Abstract: American Society of Human Genetics, 45th Annual Meeting, Minneapolis, Minnesota. *Am J Hum Genet* 57:318.
- Lubinsky MS, Adkins W, Kaveggia E (1997): Decreased maternal age with hydranencephaly. *Am J Med Genet* 69:232-234.
- Luebke HJ, Reiser CA, Pauli RM (1990): Fetal disruptions: Assessment of frequency, heterogeneity, and embryologic mechanisms in a population referred to a community-based stillbirth assessment program. *Am J Med Genet* 36:56-72.
- Mannino FL, Jones KL, Benirschke K (1977): Congenital skin defects and fetal papyraceus. *J Pediatr* 91:559-564.
- Martin ML, Khoury MJ, Cordero JF, Waters GD (1992): Trends in rates of multiple vascular disruption defects, Atlanta, 1968-1989: Is there evidence of a cocaine teratogenic epidemic? *Teratology* 45:647-653.
- Mastroiacovo P, Botto LD (1994): Chorionic villus sampling and transverse limb deficiencies: Maternal age is not a confounder. *Am J Med Genet* 53:182-186.
- Mastroiacovo P, Kallen B, Knudsen LB, Lancaster PA, Castilla EE, Mutchinick O, Robert E (1992): Absence of limbs and gross body wall defects: An epidemiological study of related rare malformation conditions. *Teratology* 46:455-464.
- Opitz JM, Kaveggia EG, Laxova R, Palister PD (1982): The diagnosis and prevention of severe mental retardation. In Hashem N, Gerald P (eds): "Proceedings, First International Conference on Preventable Aspects of Genetic Morbidity. Cairo, 1978," Vol II. pp 117-138.
- Poswillo D (1973): The pathogenesis of the first and second branchial arch syndrome. *Oral Surg* 35:302-328.
- Robins LN, Mills JL (eds) (1993): Effects of in utero exposure to street drugs. *Am J Public Health [Suppl]* 83.
- Robinson LK, Hoyme HE, Edwards DK, Jones KL (1990): Vascular pathogenesis of unilateral craniofacial defects. *J Pediatr* 111:236-239.
- Torfs CP, Velie EM, Oechsli FW, Bateson TF, Curry CJR (1994): A population-based study of gastroschisis: Demographic, pregnancy, and lifestyle risk factors. *Teratology* 50:44-53.
- Van Allen MI (1981): Fetal vascular disruptions: Mechanisms and some resulting birth defects. *Pediatr Ann* 10:219-233.
- Warkany J, Lemire RJ, Cohen MM Jr (1981): "Mental Retardation and Congenital Malformations of the Nervous System." Chicago: Year Book Medical Publishers, p 85.
- Werler MM, Mitchell AA, Shapiro S (1992a): Demographic, reproductive, medical, and environmental factors in relation to gastroschisis. *Teratology* 45:353-360.
- Werler MM, Mitchell AA, Shapiro S (1992b): First trimester maternal medication use in relation to gastroschisis. *Teratology* 45:361-367.
- Wright S (1926): Effects of age of parents on characteristics of the guinea pig. *Am Nat* 60:552-559.
- Yang P, Beaty TH, Khoury MJ, Chee E, Stewart W, Gordis L (1992): Genetic-epidemiologic study of omphalocele and gastroschisis: Evidence for heterogeneity. *Am J Med Genet* 44:668-675.